

Improving Accessibility in Books

Session 518

James Craig

Apple Accessibility Engineering

These are confidential sessions—please refrain from streaming, blogging, or taking pictures


What Is Accessibility?

And why does it matter for my readership










50 Million

Americans have a disability


U.S. Census Bureau



31 Million

Americans have a hearing impairment

S. Kochkin. Marketrak vii: Hearing loss population tops 31 million. The Hearing Review, July 2005.



10 Million

Americans have a visual impairment

American Foundation for the Blind, Public Policy Center. Statistical Snapshots, March 2008.


Accessibility on iOS

Assistive technology, built-in since iOS 3.0




“When it comes to embedding accessibility, Apple has set the standard in recent years...It is now up to other manufacturers to follow their lead.”

Robin Spinks, Royal National Institute of Blind People (RNIB), United Kingdom



“I believe the advent of accessible iBooks will be viewed by future generations as one of the landmark events in the lives of the blind.”

Bradley Hodges, American Foundation for the Blind, Access World 2010





What You Will Learn



- Basic HTML accessibility in EPUB books

What You Will Learn



- Basic HTML accessibility in EPUB books
- Basics of VoiceOver so you can test your books

What You Will Learn



- Basic HTML accessibility in EPUB books
- Basics of VoiceOver so you can test your books
- Accessibility features of iBooks Author

What You Will Learn



- Basic HTML accessibility in EPUB books
- Basics of VoiceOver so you can test your books
- Accessibility features of iBooks Author
- Techniques for accessibility in custom HTML5 widgets

Accessibility in EPUB



















optical astronomical maps, Jansky concluded that the radiation was coming from the Milky Way Galaxy and was strongest in the direction of the center of the galaxy, in the constellation of Sagittarius.



The range of frequencies in the electromagnetic spectrum that makes up the radio spectrum is very large. This means that the types of antennas that are used as radio telescopes vary widely in design, size, and configuration. At wavelengths of 30 meters to 3 meters (10 MHz to 100 MHz), they are generally either directional antenna arrays similar

to TV antennas or large stationary reflectors with moveable focal points.



Since the wavelengths being observed with these types of antennas are so long, the reflector surfaces can be constructed from coarse wire mesh. At shorter wavelengths, dish-style radio telescopes

iPad 9:41 AM

Library Sample Code Accessible Astronomy

optical astronomical maps, Jansky concluded that the radiation was coming from the Milky Way Galaxy and was strongest in the direction of the center of the galaxy, in the constellation of Sagittarius.



The range of frequencies in the electromagnetic spectrum that makes up the radio spectrum is very large. This means that the types of antennas that are used as radio telescopes vary widely in design, size, and configuration. At wavelengths of 30 meters to 3 meters (10 MHz to 100 MHz), they are generally either directional antenna arrays similar

to TV antennas or large parabolic dishes with moveable focal points.



Since the wavelengths being observed with these types of antennas are so long, the reflector surfaces can be constructed from coarse wire mesh. At shorter wavelengths, dish-style radio telescopes


4 of 14 5 of 14 9 pages left in this chapter

Full Screen OFF

iPad 9:41 AM

Library Sample Code

optical astronomical maps, Jansky concluded that the radiation was coming from the Milky Way Galaxy and was strongest in the direction of the center of the galaxy, in the constellation of Sagittarius.




The range of frequencies in the electromagnetic spectrum that makes up the radio spectrum is very large. This means that the types of antennas that are used as radio telescopes vary widely in design, size, and configuration. At wavelengths of 30 meters to 3 meters (10 MHz to 100 MHz), they are generally either directional antenna arrays similar

4 of 14

Accessible Astronomy

to TV antennas or large parabolic antennas with moveable focal points.



Since the wavelengths being observed with these types of antennas are so long, the reflector surfaces can be constructed from coarse wire mesh. At shorter wavelengths, dish-style radio telescopes

5 of 14 9 pages left in this chapter

AA Q

☀️ [Slider] ☀️

A A

Fonts

Normal

Sepia ✓


Night

Full Screen OFF

iPad 9:41 AM


Library Sample Code Accessible Astronomy

optical astronomical maps, Jansky concluded that the radiation was coming from the Milky Way Galaxy and was strongest in the direction of the center of the galaxy, in the constellation of Sagittarius.



The range of frequencies in the electromagnetic spectrum that makes up the radio spectrum is very large. This means that the types of antennas that are used as radio telescopes vary widely in design, size, and configuration. At wavelengths of 30 meters to 3 meters (10 MHz to 100 MHz), they are generally either directional antenna arrays similar

to TV antennas or large moveable focal points.



Since the wavelengths being observed with these types of antennas are so long, the reflector surfaces can be constructed from coarse wire mesh. At shorter wavelengths, dish-style radio telescopes

4 of 14 5 of 14 9 pages left in this chapter

Accessible Astronomy settings: A A, Font, Normal, Sepia, Night (checked), Full Screen OFF

optical astronomical maps, Jansky concluded that the radiation was coming from the Milky Way Galaxy and was strongest in the direction of the center of the galaxy, in the constellation of Sagittarius.



The range of frequencies in the electromagnetic spectrum that makes up the radio spectrum is very large. This means that the types of antennas that are used as radio telescopes vary widely in design, size, and configuration. At wavelengths of 30 meters to 3 meters (10 MHz to 100 MHz), they are generally either directional antenna arrays similar

to TV antennas or large stationary reflectors with moveable focal points.



Since the wavelengths being observed with these types of antennas are so long, the reflector surfaces can be constructed from coarse wire mesh. At shorter wavelengths, dish-style radio telescopes

optical astronomical maps, Jansky concluded that the radiation was coming from the Milky Way Galaxy and was strongest in the direction of the center of the galaxy, in the constellation of Sagittarius.

The range of frequencies in the electromagnetic spectrum that makes up the radio spectrum is very large. This means that the types of antennas that are used as radio telescopes vary widely in design, size, and configuration. At wavelengths of 30 meters to 3 meters (10 MHz to 100 MHz), they are generally either directional antenna arrays similar

to TV antennas or large stationary reflectors with moveable focal points.

Since the wavelengths being observed with these types of antennas are so long, the reflector surfaces can be constructed from coarse wire mesh. At shorter wavelengths, dish-style radio telescopes

Basic EPUB Accessibility

Label all content images like you would in HTML



Basic EPUB Accessibility

Label all content images like you would in HTML



```

```

Basic EPUB Accessibility

Label all content images like you would in HTML



```

```



```
 <!-- ok in some circumstances -->
```


Basic EPUB Accessibility

Label all content images like you would in HTML



```

```



```
 <!-- ok in some circumstances -->
```



```

```

Basic EPUB Accessibility



Label all content images like you would in HTML



```

```



```
 <!-- ok in some circumstances -->
```



```

```



```

```

Sample Code

optical astronomical maps, Jansky concluded that the radiation was coming from the Milky Way Galaxy and was strongest in the direction of the center of the galaxy, in the constellation of Sagittarius.

The range of frequencies in the electromagnetic spectrum that makes up the radio spectrum is very large. This means that the types of antennas that are used as radio telescopes vary widely in design, size, and configuration. At wavelengths of 30 meters to 3 meters (10 MHz to 100 MHz), they are generally either directional antenna arrays similar

Accessible Astronomy

to TV antennas or large stationary reflectors with moveable focal points.

Since the wavelengths being observed with these types of antennas are so long, the reflector surfaces can be constructed from coarse wire mesh. At shorter wavelengths, dish-style radio telescopes

optical astronomical maps, Jansky concluded that the radiation was coming from the Milky Way Galaxy and was strongest in the direction of the center of the galaxy, in the constellation of Sagittarius.

image, The Very Large Array, an interferometric array formed from many smaller radio telescopes, each dish 25 meters in diameter.

The range of frequencies in the electromagnetic spectrum that makes up the radio spectrum is very large. This means that the types of antennas that are used as radio telescopes vary widely in design, size, and configuration. At wavelengths of 30 meters to 3 meters (10 MHz to 100 MHz), they are generally either directional antenna arrays similar

to TV antennas or large stationary reflectors with moveable focal points.

image, Combined x-ray, microwave, and visible spectrum photograph showing elliptical galaxy Centaurus A, with radio-emitting jets and lobes emanating from its central black hole.


Since the wavelengths being observed with these types of antennas are so long, the reflector surfaces can be constructed from coarse wire mesh. At shorter wavelengths, dish-style radio telescopes

optical astronomical maps, Jansky concluded that the radiation was coming from the Milky Way Galaxy and was strongest in the direction of the center of the galaxy, in the constellation of Sagittarius.



The range of frequencies in the electromagnetic spectrum that makes up the radio spectrum is very large. This means that the types of antennas that are used as radio telescopes vary widely in design, size, and configuration. At wavelengths of 30 meters to 3 meters (10 MHz to 100 MHz), they are generally either directional antenna arrays similar

to TV antennas or large stationary reflectors with moveable focal points.



Since the wavelengths being observed with these types of antennas are so long, the reflector surfaces can be constructed from coarse wire mesh. At shorter wavelengths, dish-style radio telescopes

optical astronomical maps, Jansky concluded that the radiation was coming from the Milky Way Galaxy and was strongest in the direction of the center of the galaxy, in the constellation of Sagittarius.



The range of frequencies in the electromagnetic spectrum that makes up the radio spectrum is very large. This means that the types of antennas that are used as radio telescopes vary widely in design, size, and configuration. At wavelengths of 30 meters to 3 meters (10 MHz to 100 MHz), they are generally either directional antenna arrays similar

to TV antennas or large stationary reflectors with moveable focal points.



Since the wavelengths being observed with these types of antennas are so long, the reflector surfaces can be constructed from coarse wire mesh. At shorter wavelengths, dish-style radio telescopes

Demo

Using VoiceOver in iOS and iBooks

Jason Barry

Developer Publications

Review

What we have learned so far about EPUB accessibility

Review

What we have learned so far about EPUB accessibility

- View your images in different contexts and different themes

Review

What we have learned so far about EPUB accessibility


- View your images in different contexts and different themes
- Label all your content images as you would in HTML

Review

What we have learned so far about EPUB accessibility

- View your images in different contexts and different themes
- Label all your content images as you would in HTML
- Test your books with VoiceOver!


Accessibility in iBooks Author




CHAPTER 1

Welcome

- 1.1 Molecular Pageant
- 1.2 Cells Are the Unit of Life
- 1.3 Small Creatures Rule the Earth
- 1.4 Green Planet
- 1.5 Project-Based Learning





Movie Chemical Language of Pheromones

Ants communicate using combinations of ten to twenty chemical signals.

In July and August of 2011 the team filmed in Gorongosa National Park in the southern African nation of Mozambique, bringing home feature material for four chapters in ecology. While shooting in the rain forest atop Gorongosa Mountain, we spent a little time collecting. The carpenter ant to the left, collected on our trip, is a species of the genus *Camponotus* that was previously unknown to science.




Our goal is to have students see the living world the way a naturalist sees it.

To “see,” for example, the chemical environment of organisms, such as the plumes of **pheromone** and territory-marking molecules that organisms use to communicate. Ants have been the study of a lifetime for Edward O. Wilson. We’ll bring some special lessons to students from the world of ants.

Interactive Insect Body Plan

The Insect Body Plan



The word insect is derived from the Greek meaning “cut into sections.” The evolutionary success of insects and their distinctive physiology can be measured by this: as much as 90 percent of all animal species are insects.

Enhancing Accessibility

Considerations for persons with disabilities

Adjustable Font Sizes

Color Concerns

Captions and Subtitles

Descriptive Labels

Enhancing Accessibility

Considerations for persons with disabilities

Adjustable Font Sizes

Color Concerns

Captions and Subtitles

Descriptive Labels

Section 1

Interstellar Time Travel

FACTS

1. Gravity is the curvature of space-time
2. Space is awesome
3. We're in space and space is the place



The Milky Way is the galaxy that contains the Earth. This name derives from its appearance as a dim "milky" glowing band arching across the night sky, in which the naked eye cannot distinguish individual stars. The Milky Way appears like a band because it is a disk-shaped structure being viewed from inside. The fact that this faint band of light is made up of stars was proven in 1610 when Galileo Galilei used his telescope to resolve it into individual stars. In the 1920s, observations by astronomer Edwin Hubble showed that the Milky Way is just one of many galaxies.

The Milky Way is a barred spiral galaxy 100,000–120,000 light-years in diameter containing 200–400 billion stars. It may contain at least as many planets, with 10 billion of those orbiting in the habitable zone of their parent stars. The Solar System is located within the disk, around two thirds of the way out from the Galactic Center, on the inner edge of a spiral-shaped concentration of gas and dust called the Orion–Cygnus Arm. The stars in the inner 10,000 light-years are organized in a bulge and one or more bars. The very center is marked by an intense radio source named Sagittarius A which is likely to be a supermassive black hole. The Galaxy rotates once every 15 to 50 million years. The Galaxy as a whole is moving at a velocity of 552 to 630 km per second, depending on the relative frame of reference. It is estimated to be about 13.2 billion years old, nearly as old as the Universe. Surrounded by several smaller satellite galaxies, the Milky

Way is part of the Local Group of galaxies, which forms a subcomponent of the Virgo Supercluster.

When observing the night sky, the term "Milky Way" is limited to the hazy band of white light some 30 degrees wide arcing across the sky (although all of the stars that can be seen with the naked eye are part of the Milky Way Galaxy). The light in this band originates from un-resolved stars and other material that lie within the Galactic plane. Dark regions within the band, such as the Great Rift and the Coalsack, correspond to areas where light from distant stars is blocked by interstellar dust.

The Milky Way has a relatively low surface brightness. Its visibility can be greatly reduced by background light such as light pollution or stray light

from the moon. It is readily visible when the limiting magnitude is +5.1 or better, while showing a great deal of detail at +6.1. This makes the Milky Way difficult to see from any brightly-lit urban or

suburban location but very prominent when viewed from a rural area when the moon is below the horizon.

The Milky Way passes through parts of roughly 30 constellations. The center of the Galaxy lies in the direction of the constellation Sagittarius; it is here that the Milky Way is brightest. From Sagittarius, the hazy band of white light appears to pass

westward to the Galactic anticenter in Auriga. The band then continues westward the rest of the way around the sky back to Sagittarius. The fact that the band divides the night sky into two roughly equal hemispheres indicates that the Solar System lies close to the Galactic plane. The Galactic plane is inclined by about 60 degrees to the ecliptic (the plane of the Earth's orbit). Relative to the celestial equator, it

passes as far north as the constellation of Cassiopeia and as far south as the constellation of Crux, indicating the high inclination of Earth's equatorial plane and the plane of the ecliptic relative to

Movie 1.1 Final launch of the Space Shuttle Atlantis



Launch of the Space Shuttle Atlantis. Enable closed caption display in Video settings.

diameter containing 200-400 billion stars. It may contain at least as many planets, with 10 billion of those orbiting in the habitable zone of their parent stars. The Solar System is located within the disk, around two thirds of the way out from the Galactic Center, on the inner edge of a spiral-shaped concentration of gas and dust called the Orion-Cygnus Arm. The stars in the inner 10,000 light-years are organized in a bulge and one or more bars. The very center is marked by an intense radio source named Sagittarius A which is likely to be a supermassive black hole. The Galaxy rotates once every 15 to 50 million years. The Galaxy as a whole is moving at a velocity of 552 to 630 km per second, depending on the relative frame of reference. It is estimated to be about 13.2 billion years old, nearly as old as the Universe. Surrounded by several smaller satellite galaxies, the Milky Way is part of the Local Group of galaxies, which forms a subcomponent of the Virgo Supercluster.




Movie 1.1 Final launch of the Space Shuttle...

When observing the night sky, the term "Milky Way" is limited to the hazy band of white light some 30 degrees wide arcing across the sky (although all of the stars that can be seen with the naked eye are part of the Milky Way Galaxy). The light in this band originates from un-resolved stars and other material that lie within the Galactic plane. Dark regions within the band, such as the Great Rift and the Coalsack, correspond to areas where light from distant stars is blocked by interstellar dust.

The Milky Way has a relatively low surface brightness. Its visibility can be greatly reduced by background light such as light pollution or stray light from the moon. It is readily visible when the limiting magnitude is +5.1 or better, while showing a great deal of detail at +6.1. This makes the Milky Way difficult to see from any brightly-lit urban or suburban location but very prominent when viewed from a rural area when the moon is below the horizon.

The Milky Way passes through parts of roughly 30 constellations. The center of the Galaxy lies in the direction of the constellation Sagittarius; it is here that the Milky Way is brightest. From Sagittarius, the hazy band of white light appears to pass westward to the Galactic anticenter in Auriga. The band then continues westward the rest of the way around the sky back to Sagittarius. The fact that the band divides the night sky into two roughly equal hemispheres indicates that the Solar System lies close to the Galactic plane. The Galactic plane is inclined by about 60 degrees to the ecliptic (the plane of the Earth's orbit). Relative to the celestial equator, it passes as far north as the constellation of Cassiopeia and as far south as the constellation of Crux, indicating the high inclination of Earth's equatorial plane and the plane of the ecliptic relative to the Galactic plane. The north Galactic pole is



stars. The Solar System is located with way out from the Galactic Center, on the concentration of gas and dust called the inner 10,000 light-years are organized very center is marked by an intense radiation is likely to be a supermassive black hole to 50 million years. The Galaxy as a whole is moving at a velocity of 552 to 630 km per second, depending on the relative frame of reference. It is estimated to be about 13.2 billion years old, nearly as old as the Universe. Surrounded by several smaller satellite galaxies, the Milky Way is part of the Local Group of galaxies, which forms a subcomponent of the Virgo Supercluster.




Movie 1.1 Final launch of the Space Shuttle...

When observing the night sky, the term "Milky Way" is limited to the hazy band of white light some 30 degrees wide arcing across the sky (although all of the stars that can be seen with the naked eye are part of the Milky Way Galaxy). The light in this band originates from un-resolved stars and other material that lie within the Galactic plane. Dark regions within the band, such as the Great Rift and the Coalsack, correspond to areas where light from distant stars is blocked by interstellar dust.

The Milky Way has a relatively low surface brightness. Its visibility can be greatly reduced by background light such as light pollution or stray light from the moon. It is readily visible when the limiting magnitude is +5.1 or better, while showing a great deal of detail at +6.1. This makes the Milky Way difficult to see from any brightly-lit urban or suburban location but very prominent when viewed from a rural area when the moon is below the horizon.

The Milky Way passes through parts of roughly 30 constellations. The center of the Galaxy lies in the direction of the constellation Sagittarius; it is here that the Milky Way is brightest. From Sagittarius, the hazy band of white light appears to pass westward to the Galactic anticenter in Auriga. The band then continues westward the rest of the way around the sky back to Sagittarius. The fact that the band divides the night sky into two roughly equal hemispheres indicates that the Solar System lies close to the Galactic plane. The Galactic plane is inclined by about 60 degrees to the ecliptic (the plane of the Earth's orbit). Relative to the celestial equator, it passes as far north as the constellation of Cassiopeia and as far south as the constellation of Crux, indicating the high inclination of Earth's equatorial plane and the plane of the ecliptic relative to the Galactic plane. The north Galactic pole is

Text adjustment menu with a slider and two 'A' buttons.



stars. The Solar System is located with way out from the Galactic Center, on the concentration of gas and dust called the inner 10,000 light-years are organized very center is marked by an intense radiation is likely to be a supermassive black hole to 50 million years. The Galaxy as a whole is moving at a velocity of 552 to 630 km per second, depending on the relative frame of reference. It is estimated to be about 13.2 billion years old, nearly as old as the Universe. Surrounded by several smaller satellite galaxies, the Milky Way is part of the Local Group of galaxies, which forms a subcomponent of the Virgo Supercluster.



Movie 1.1 Final launch of the Space Shuttle...

When observing the night sky, the term "Milky Way" is limited to the hazy band of white light some 30 degrees wide arcing across the sky (although all of the stars that can be seen with the naked eye are part of the Milky Way Galaxy). The light in this band originates from un-resolved stars and other material that lie within the Galactic plane. Dark regions within the band, such as the Great Rift and the Coalsack, correspond to areas where light from distant stars is blocked by interstellar dust.

The Milky Way has a relatively low surface brightness. Its visibility can be greatly reduced by background light such as light pollution or stray light from the moon. It is readily visible when the limiting magnitude is +5.1 or better, while showing a great deal of detail at +6.1. This makes the Milky Way difficult to see from any brightly-lit urban or suburban location but very prominent when viewed from a rural area when the moon is below the horizon.

The Milky Way passes through parts of roughly 30 constellations. The center of the Galaxy lies in the direction of the constellation Sagittarius; it is here that the Milky Way is brightest. From Sagittarius, the hazy band of white light appears to pass westward to the Galactic anticenter in Auriga. The band then continues westward the rest of the way around the sky back to Sagittarius. The fact that the band divides the night sky into two roughly equal hemispheres indicates that the Solar System lies close to the Galactic plane. The Galactic plane is inclined by about 60 degrees to the ecliptic (the plane of the Earth's orbit). Relative to the celestial equator, it passes as far north as the constellation of Cassiopeia and as far south as the constellation of Crux, indicating the high inclination of Earth's equatorial plane and the plane of the ecliptic relative to the Galactic plane. The north Galactic pole is

rotates once every 15 to 50 million years. The Galaxy as a whole is moving at a velocity of 552 to 630 km per second, depending on the relative frame of reference. It is estimated to be about 13.2 billion years old, nearly as old as the Universe. Surrounded by several smaller satellite galaxies, the Milky Way is part of the Local Group of galaxies, which forms a subcomponent of the Virgo Supercluster.



Movie 1.1 Final launch of the Space Shuttle...

When observing the night sky, the term "Milky Way" is limited to the hazy band of white light some 30 degrees wide arcing across the sky (although all of the stars that can be seen with the naked eye are part of the Milky Way Galaxy). The light in this band originates from un-resolved stars and other material that lie within the Galactic plane. Dark regions within the band, such as the Great Rift and the Coalsack, correspond to areas where light from distant stars is blocked by interstellar dust.

The Milky Way has a relatively low surface

Enhancing Accessibility

Considerations for persons with disabilities

Adjustable Font Sizes

Color Concerns

Captions and Subtitles

Descriptive Labels

Enhancing Accessibility

Considerations for persons with disabilities

Adjustable Font Sizes


Color Concerns

Captions and Subtitles

Descriptive Labels


Color-Blindness

Example of what not to do




Color-Blindness

Do not convey meaning by color alone



Color-Blindness


Do not convey meaning by color alone



- Not color-blind (92%)
- Color-blind (8%)

Color-Blindness

Do not convey meaning by color alone



- Not color-blind (92%)
- Color-blind (8%)

Not color-blind (92%)
Color-blind (8%)

SECTION 4

Review

Summary

Ecology is the scientific study of the relations between organisms and their environments. One goal of ecology is to discover the rules that explain the patterns we see in the natural world. Fundamental to these investigations is assessing the influence of biotic and abiotic factors.


Populations of organisms that share a habitat and interact with one another form a community. On a larger scale, the interactions and interdependencies that connect organisms and their environments comprise an ecosystem.

Biomes are communities on a very large scale, identified by their characteristic climate and prevailing vegetation. At the highest level, terrestrial biomes include forest, grassland, desert, and tundra, and aquatic biomes are divided between freshwater and marine. There are numerous finer distinctions, based on patterns found in specific climates and regions of the world.

Review 1

Question 5 of 5

Match the biomes on this map of the Americas:

[Try Again](#)

SECTION 4

Review

Summary

Ecology is the scientific study of the relations between organisms and their environments. One goal of ecology is to discover the rules that explain the patterns we see in the natural world. Fundamental to these investigations is assessing the influence of biotic and abiotic factors.

Populations of organisms that share a habitat and interact with one another form a community. On a larger scale, the interactions and interdependencies that connect organisms and their environments comprise an ecosystem.

Biomes are communities on a very large scale, identified by their characteristic climate and prevailing vegetation. At the highest level, terrestrial biomes include forest, grassland, desert, and tundra, and aquatic biomes are divided between freshwater and marine. There are numerous finer distinctions, based on patterns found in specific climates and regions of the world.

Review 1

Question 5 of 5

Match the biomes on this map of the Americas:

[Try Again](#)

SECTION 4

Review

Summary

Ecology is the scientific study of the relations between organisms and their environments. One goal of ecology is to discover the rules that explain the patterns we see in the natural world. Fundamental to these investigations is assessing the influence of biotic and abiotic factors.

Populations of organisms that share a habitat and interact with one another form a community. On a larger scale, the interactions and interdependencies that connect organisms and their environments comprise an ecosystem.

Biomes are communities on a very large scale, identified by their characteristic climate and prevailing vegetation. At the highest level, terrestrial biomes include forest, grassland, desert, and tundra, and aquatic biomes are divided between freshwater and marine. There are numerous finer distinctions, based on patterns found in specific climates and regions of the world.

Review 1



Enhancing Accessibility

Considerations for persons with disabilities

Adjustable Font Sizes

Color Concerns

Captions and Subtitles

Descriptive Labels

Enhancing Accessibility


Considerations for persons with disabilities


Adjustable Font Sizes

Color Concerns

Captions and Subtitles

Descriptive Labels





Enhancing Accessibility

Considerations for persons with disabilities

Adjustable Font Sizes

Color Concerns

Captions and Subtitles

Descriptive Labels

Enhancing Accessibility


Considerations for persons with disabilities


Adjustable Font Sizes

Color Concerns

Captions and Subtitles

Descriptive Labels






Widget

Layout Interaction

Layout




Title
Label

Caption

Background
Margin

Accessibility Description

Wide shot of Jupiter, with swirling gas clouds, including the Great Red Spot, a storm so large it dwarfs our own Earth



Widget

Layout Interaction

Layout

Title
Label

Caption

Background
Margin

Accessibility Description

Wide shot of Jupiter, with swirling gas clouds, including the Great Red Spot, a storm so large it dwarfs our own Earth

the Galactic plane. The north Galactic pole is situated at right ascension 12h 51m 06s, declination +90°. The south Galactic pole is near beta Comae Berenices, at right ascension 13h 52m 54s, declination -90°. Sculptor's Belt is a region of stars near alpha Centauri, extending on the time of night from the southern horizon to the north. It can appear relatively low in the sky, but it is not visible from servers from about 65 degrees north. The Earth's surface the Milky Way is visible twice a day. The Milky Way stars and may have up to 100 billion stars, depending on which depends on the number of stars in the galaxy, which are hard to detect, especially those within 100 ly (90 pc) from the Sun. The Andromeda Galaxy contains about 1 trillion stars. Filling the space between stars is the interstellar medium, the gas and dust between stars. The extent in radius of the Milky Way is about 150,000 light years. The layer ranges from the center to the edge. There are thousands of stars in the Milky Way. Microlensing indicates that there may be many more stars in the galaxy than we can see. There are stars in the galaxy that are much smaller than the Sun, and some indicate that there may be more stars than the Sun.

Widget

Layout Interaction

Full-screen only

Questions	# of Answers
1 - Planets in the Milky Way	4

+ -

Accessibility Description

Question

Match the planets to their location in the solar system.

Match the images of these planets to their corresponding positions in our solar system.

Planets in the Milky Way

Check Answer

the Galactic plane. The north Galactic pole is situated at right ascension 12h 51m 06s, declination +90°. The south Galactic pole is near beta Comae Berenices, at right ascension 13h 02m 48s, declination -90°. Sculptor's Belt is a region of stars near alpha Centauri, extending on the time of night. It can appear as a band of stars from the Earth's surface. The Milky Way is about 65 degrees wide. The Milky Way's surface may have up to 100 billion stars, depending on the number of stars which are hard to detect, especially those 100 ly (90 pc) from the Sun. The Andromeda Galaxy contains about 2 trillion stars. The space between stars is vast. The interstellar medium is comparable in extent to the layer of the atmosphere. The interstellar layer ranges from a few parsecs to thousands of parsecs. Gravitational microlensing indicates that there may be billions of stars in the Galaxy. Measurements indicate that the Milky Way is surrounded by billions of stars, more than the solar system.

Widget

Layout Interaction

Full-screen only

Questions	# of Answers
1 - Planets in the Milky Way	4

Accessibility Description

Question

Match the planets to their location in the solar system.

Match the images of these planets to their corresponding positions in our solar system.

Planets in the Milky Way

Check Answer

Demo

Adding accessibility to content in iBooks Author

Review

What we have learned so far about iBooks accessibility

Review

What we have learned so far about iBooks accessibility

- Font size adjustment is already built-in

Review

What we have learned so far about iBooks accessibility

- Font size adjustment is already built-in
- Don't convey meaning with color alone

Review

What we have learned so far about iBooks accessibility

- Font size adjustment is already built-in
- Don't convey meaning with color alone
- Add captions or subtitles to video where appropriate

Review

What we have learned so far about iBooks accessibility

- Font size adjustment is already built-in
- Don't convey meaning with color alone
- Add captions or subtitles to video where appropriate
- Add alternative text for images, widgets, etc.

Review

What we have learned so far about iBooks accessibility

- Font size adjustment is already built-in
- Don't convey meaning with color alone
- Add captions or subtitles to video where appropriate
- Add alternative text for images, widgets, etc.
- Test your books with VoiceOver!

Custom HTML Widgets





Key	Value
+ Property List	17 Items
Localization	English
Application-Group	Developer Tools
Executable	Property List Editor
BundleIdentifier	com.apple.PropertyListEditor
BundleStyle	APPL
DictionaryVersion	6.0
PrincipalClass	NSApplication

PLIST

“ARIA allows web pages, or portions of web pages, to declare themselves as applications...”

WAI-ARIA 1.0 Candidate Recommendation, 18 January 2011
<http://w3.org/TR/wai-aria/>

Using WAI-ARIA

Accessible Rich Internet Applications

Using WAI-ARIA

Accessible Rich Internet Applications

- Assign semantic roles

Using WAI-ARIA

Accessible Rich Internet Applications

- Assign semantic roles



Using WAI-ARIA

Accessible Rich Internet Applications

- Assign semantic roles


```
<div class="front" role="button">  
</div>
```



Using WAI-ARIA

Accessible Rich Internet Applications

- Assign semantic roles




Using WAI-ARIA

Accessible Rich Internet Applications

- Assign semantic roles

```
<div class="back" role="dialog">  
</div>
```



Using WAI-ARIA

Accessible Rich Internet Applications

- Assign semantic roles
- Update states and properties

Using WAI-ARIA

Accessible Rich Internet Applications

- Assign semantic roles
- Update states and properties



Using WAI-ARIA

Accessible Rich Internet Applications

- Assign semantic roles
- Update states and properties

```
<div class="front" role="button"  
  aria-label="The Moon">  
</div>
```



Using WAI-ARIA

Accessible Rich Internet Applications

- Assign semantic roles
- Update states and properties

```
<div class="front" role="button"  
  aria-label="The Moon">  
</div>
```




Using WAI-ARIA

Accessible Rich Internet Applications

- Assign semantic roles
- Update states and properties

```
<div class="front" role="button"  
  aria-label="The Moon">  
</div>
```


```
<div class="back" role="dialog"  
  aria-hidden="true">  
</div>
```



Using WAI-ARIA

Accessible Rich Internet Applications

- Assign semantic roles
- Update states and properties




Using WAI-ARIA

Accessible Rich Internet Applications

- Assign semantic roles
- Update states and properties

```
// show the dialog to VO  
dialog.setAttribute  
  ('aria-hidden', 'false');
```




Using WAI-ARIA

Accessible Rich Internet Applications

- Assign semantic roles
- Update states and properties

```
// show the dialog to VO  
dialog.setAttribute  
  ('aria-hidden', 'false');
```




Using WAI-ARIA

Accessible Rich Internet Applications

- Assign semantic roles
- Update states and properties

```
// show the dialog to VO
dialog.setAttribute
  ('aria-hidden', 'false');

// hide the buttons from VO
buttons.setAttribute
  ('aria-hidden', 'true');
```



Using WAI-ARIA

Accessible Rich Internet Applications

- Assign semantic roles
- Update states and properties
- Manage focus

Using WAI-ARIA

Accessible Rich Internet Applications

- Assign semantic roles
- Update states and properties
- Manage focus



Using WAI-ARIA

Accessible Rich Internet Applications


- Assign semantic roles
- Update states and properties
- Manage focus



Using WAI-ARIA

Accessible Rich Internet Applications

- Assign semantic roles
- Update states and properties
- Manage focus




Using WAI-ARIA

Accessible Rich Internet Applications

- Assign semantic roles
- Update states and properties
- Manage focus

```
firstImageInDialog.focus();
```



Using WAI-ARIA

Accessible Rich Internet Applications

- Assign semantic roles
- Update states and properties
- Manage focus



Using WAI-ARIA

Accessible Rich Internet Applications

- Assign semantic roles
- Update states and properties
- Manage focus

```
triggerButton.focus();
```



Using WAI-ARIA

Accessible Rich Internet Applications

Using WAI-ARIA

Accessible Rich Internet Applications

- Retrofit old content without gutting it

Using WAI-ARIA

Accessible Rich Internet Applications

- Retrofit old content without gutting it

```
<div class="front" role="button" aria-label="The Moon"></div>
```

Using WAI-ARIA

Accessible Rich Internet Applications

- Retrofit old content without gutting it

```
<div class="front" role="button" aria-label="The Moon"></div>  
el.setAttribute('role', 'button');
```

Using WAI-ARIA

Accessible Rich Internet Applications

- Retrofit old content without gutting it

```
<div class="front" role="button" aria-label="The Moon"></div>  
el.setAttribute('role', 'button');  
el.setAttribute('aria-label', 'The Moon');
```

Using WAI-ARIA

Accessible Rich Internet Applications

- Retrofit old content without gutting it

```
<div class="front" role="button" aria-label="The Moon"></div>  
el.setAttribute('role', 'button');  
el.setAttribute('aria-label', 'The Moon');
```

- Update your application state when you change element style

Using WAI-ARIA

Accessible Rich Internet Applications

- Retrofit old content without gutting it

```
<div class="front" role="button" aria-label="The Moon"></div>  
el.setAttribute('role', 'button');  
el.setAttribute('aria-label', 'The Moon');
```

- Update your application state when you change element style

```
el.className = 'hidden';
```

Using WAI-ARIA

Accessible Rich Internet Applications

- Retrofit old content without gutting it

```
<div class="front" role="button" aria-label="The Moon"></div>  
el.setAttribute('role', 'button');  
el.setAttribute('aria-label', 'The Moon');
```

- Update your application state when you change element style

```
el.className = 'hidden';  
el.setAttribute('aria-hidden', 'true');
```

Focus Management

Setting VoiceOver cursor focus on iOS

Focus Management

Setting VoiceOver cursor focus on iOS

- VoiceOver cursor focus is not the same as keyboard focus!

Focus Management

Setting VoiceOver cursor focus on iOS

- VoiceOver cursor focus is not the same as keyboard focus!
- Don't focus hidden elements (and don't hide elements that have focus)

Focus Management

Setting VoiceOver cursor focus on iOS

- VoiceOver cursor focus is not the same as keyboard focus!
- Don't focus hidden elements (and don't hide elements that have focus)

```
el.removeAttribute('aria-hidden'); // show the element...
```


Focus Management

Setting VoiceOver cursor focus on iOS

- VoiceOver cursor focus is not the same as keyboard focus!
- Don't focus hidden elements (and don't hide elements that have focus)

```
el.removeAttribute('aria-hidden'); // show the element...  
el.focus(); // ...before you move focus to it
```

- Use tabindex if needed

Focus Management

Setting VoiceOver cursor focus on iOS

- VoiceOver cursor focus is not the same as keyboard focus!
- Don't focus hidden elements (and don't hide elements that have focus)

```
el.removeAttribute('aria-hidden'); // show the element...  
el.focus(); // ...before you move focus to it
```

- Use tabIndex if needed

```
el.tabIndex = 0; // make sure the element is focusable first...
```

Focus Management

Setting VoiceOver cursor focus on iOS

- VoiceOver cursor focus is not the same as keyboard focus!
- Don't focus hidden elements (and don't hide elements that have focus)

```
el.removeAttribute('aria-hidden'); // show the element...  
el.focus(); // ...before you move focus to it
```

- Use tabIndex if needed

```
el.tabIndex = 0; // make sure the element is focusable first...  
el.focus(); // ...or it will throw an error in some contexts
```

Demo

Creating a custom accessible widget

Review

What we have learned so far about HTML widget accessibility

Review

What we have learned so far about HTML widget accessibility

- Standard HTML accessibility techniques still apply

Review

What we have learned so far about HTML widget accessibility

- Standard HTML accessibility techniques still apply
- Use ARIA to declare application semantics where HTML cannot

Review

What we have learned so far about HTML widget accessibility

- Standard HTML accessibility techniques still apply
- Use ARIA to declare application semantics where HTML cannot
- Manage focus when application state changes

Review

What we have learned so far about HTML widget accessibility

- Standard HTML accessibility techniques still apply
- Use ARIA to declare application semantics where HTML cannot
- Manage focus when application state changes
- Test your books with VoiceOver!



Takeaways



- Books are for everyone

Takeaways



- Books are for everyone
- Accessibility is for everyone, not just those with disabilities

Takeaways



- Books are for everyone
- Accessibility is for everyone, not just those with disabilities
- Adding accessibility to books is easy

Takeaways



- Books are for everyone
- Accessibility is for everyone, not just those with disabilities
- Adding accessibility to books is easy
- Test your books with VoiceOver!

More Information

Vicki Murley

Safari Technologies Evangelist

vicki@apple.com

Apple Accessibility

<http://www.apple.com/accessibility>

WAI-ARIA

<http://www.w3.org/TR/wai-aria>

Apple Developer Forums

<http://devforums.apple.com>

Related Sessions

Accessibility for OS X

Marina
Tuesday 10:15AM

Building Books with iBooks Author

Mission
Tuesday 11:30AM

Accessibility for iOS

Russian Hill
Wednesday 9:00AM

HTML, CSS, and DOM for Book Authors

Nob Hill
Wednesday 3:15PM

 **WWDC2012**